

A Fitting Formula for the Non-Linear Evolution of the Bispectrum

Román Scoccimarro¹ and H.M.P. Couchman²

¹*Institute for Advanced Study, School of Natural Sciences, Einstein Drive, Princeton, NJ 08540, USA*

²*Department of Physics and Astronomy, McMaster University, Hamilton, Ontario, L8S 4M1, Canada*

1 February 2008

ABSTRACT

We present a fitting formula for the non-linear evolution of the bispectrum in CDM models, obtained from measurements in high resolution numerical simulations. The formula interpolates between the perturbative and highly non-linear regimes, and generalizes previous results obtained for scale-free initial conditions.

Key words: cosmology: large-scale structure of the universe; methods: numerical

1 INTRODUCTION

The large-scale structure of the universe observed in galaxy surveys is thought to arise from the growth of primordial fluctuations due to gravitational instability. Making accurate predictions of gravitational clustering is essential if we are to compare theories with observations and put constraints on cosmological parameters.

Gravitational clustering is a non-linear process: when fluctuations are sufficiently small the equations of motion can be solved perturbatively, but at small enough scales one has resort to numerical simulations to accurately follow the dynamics. Non-linearity of the equations of motion implies that the mass distribution becomes non-Gaussian, even if the initial conditions are Gaussian (Peebles 1980, Fry 1984).

The power spectrum, or two-point correlation function, has become the most common statistical measure of gravitational clustering. Its importance rests on the fact that for a Gaussian random field, the power spectrum completely characterizes its statistical properties. Hamilton et al. (1991) suggested a useful way of thinking about the non-linear evolution of the power spectrum, summarized by a simple fitting formula obtained from numerical simulations. The accuracy of this procedure has been improved (e.g. Jain, Mo & White 1995; Peacock & Dodds 1996) and extensively used in the literature to predict statistical properties related to the density power spectrum.

A complete description of the non-Gaussian distribution that results from gravitational instability, however, requires knowledge of higher-order correlation functions. A number of useful statistics have emerged in the last decade to become standard probes of non-Gaussianity in large-scale structure. These include the higher-order moments of the smoothed density field (e.g. skewness and kurtosis; Peebles 1980; Juszkiewicz, Colombi & Bouchet 1993; Bernardeau

1994) and the three-point correlation function in Fourier space, the bispectrum (Fry 1984; Scoccimarro et al. 1998, hereafter SCFFHM) or in real space (Jing & Börner 1998; Frieman & Gaztañaga 1999).

A fitting formula for the non-linear evolution of one-point higher-order moments is given by extended perturbation theory (EPT, Colombi et al. 1997). The EPT prescription uses the perturbative expressions for skewness, kurtosis, and so on, but with an “effective” spectral index that becomes a function of the variance σ^2 of the smoothed density field. This function is obtained numerically by fitting to measurements of the density field moments in N-body simulations. The EPT prescription is able to fit reasonably well the non-linear evolution of higher-order moments and thus the one-point probability distribution function with this single function obtained from numerical simulations.

Probing non-Gaussianity beyond one-point statistics gives precious additional information that can be used to learn about cosmology. The bispectrum is the lowest-order statistic that probes the shape of large-scale structures generated by gravitational clustering. Measurements of the bispectrum in galaxy redshift catalogs have been proposed as a way to constrain galaxy biasing (Fry 1994; Hivon et al. 1995; Matarrese, Verde & Heavens 1997) and thus break the degeneracies present in power spectrum measurements between the matter density of the universe Ω_m and galaxy bias. Such measurements have recently been done in IRAS redshift surveys (Scoccimarro et al. 2000; Feldman et al. 2000), providing constraints on galaxy bias, Ω_m and non-Gaussian primordial fluctuations. In addition, upcoming large redshift surveys such as 2dF and SDSS will determine the bispectrum with unprecedented accuracy.

In this work we study the description of the non-linear evolution of the bispectrum. A fitting formula for the bispectrum was proposed by Scoccimarro & Frieman (1999),

arXiv:astro-ph/0009427v1 26 Sep 2000

hereafter SF, that interpolates between perturbative and the highly non-linear regime, assuming scale-free initial conditions. Since current models for structure formation are not scale-free, these results could not be used to make realistic predictions to be compared with observations. In this work, we extend the SF results to provide a fitting formula for the non-linear evolution of the bispectrum in CDM models.

2 THE BISPECTRUM

We denote by $\tilde{\delta}$ the Fourier transform of the density contrast δ obtained from the density field $\rho = \bar{\rho}(1 + \delta)$, where $\bar{\rho}$ is the mean density. The power spectrum $P(k)$ and the bispectrum $B_{123} = B(k_1, k_2, k_3)$ are defined as

$$\begin{aligned} \langle \tilde{\delta}(\mathbf{k}_1) \tilde{\delta}(\mathbf{k}_2) \rangle &= \delta_D(\mathbf{k}_1 + \mathbf{k}_2) P(k) \\ \langle \tilde{\delta}(\mathbf{k}_1) \tilde{\delta}(\mathbf{k}_2) \tilde{\delta}(\mathbf{k}_3) \rangle &= \delta_D(\mathbf{k}_1 + \mathbf{k}_2 + \mathbf{k}_3) B_{123}. \end{aligned} \quad (1)$$

In this paper, we assume Gaussian initial conditions. In this case, the bispectrum generated by gravitational instability at large scales is given by tree-level (second-order) perturbation theory, and reads (Fry 1984)

$$B_{123} = 2 F_2(\mathbf{k}_1, \mathbf{k}_2) P_1 P_2 + \text{cyc.}, \quad (2)$$

where $P_i \equiv P(k_i)$, and

$$F_2(\mathbf{k}_1, \mathbf{k}_2) = \frac{5}{7} + \frac{1}{2} \frac{\mathbf{k}_1 \cdot \mathbf{k}_2}{k_1 k_2} \left(\frac{k_1}{k_2} + \frac{k_2}{k_1} \right) + \frac{2}{7} \left(\frac{\mathbf{k}_1 \cdot \mathbf{k}_2}{k_1 k_2} \right)^2. \quad (3)$$

Note that Eq.(1) implies that the bispectrum is defined only for closed triangles formed by the wave vectors $\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3$. It is convenient to define the reduced bispectrum Q_{123} as

$$Q_{123} = \frac{B_{123}}{P_1 P_2 + P_2 P_3 + P_3 P_1}, \quad (4)$$

which takes away most of the dependence on scale and cosmology.

SF gave a fitting formula for the non-linear evolution of the bispectrum from scale-free initial conditions. They replaced the kernel in Eq. (3) by the effective kernel

$$\begin{aligned} F_2^{\text{eff}}(\mathbf{k}_1, \mathbf{k}_2) &= \frac{5}{7} a(n, k_1) a(n, k_2) \\ &+ \frac{1}{2} \frac{\mathbf{k}_1 \cdot \mathbf{k}_2}{k_1 k_2} \left(\frac{k_1}{k_2} + \frac{k_2}{k_1} \right) b(n, k_1) b(n, k_2) \\ &+ \frac{2}{7} \left(\frac{\mathbf{k}_1 \cdot \mathbf{k}_2}{k_1 k_2} \right)^2 c(n, k_1) c(n, k_2), \end{aligned} \quad (5)$$

where the functions $a(n, k)$, $b(n, k)$, and $c(n, k)$ are chosen to interpolate between these two regimes according to the one-loop PT and N-body results (SCFFHM). This yields $(-2 \leq n \leq 0)$

$$a(n, k) = \frac{1 + [0.7 Q_3(n)]^{1/2} (k R_0)^{n+6}}{1 + (k R_0)^{n+6}}, \quad (6)$$

$$b(n, k) = \frac{1 + 0.2 (n+3) (k R_0)^{n+3}}{1 + (k R_0)^{n+3.5}}, \quad (7)$$

$$c(n, k) = \frac{1 + 4.5/[1.5 + (n+3)^4] (k R_0)^{n+3}}{1 + (k R_0)^{n+3.5}}, \quad (8)$$

where R_0 is a measure of the non-linear scale, defined from

Table 1. Numerical Simulations Parameters

Model	z	$h(z)$	$\Omega_m(z)$	$\Omega_\Lambda(z)$	Γ	$\sigma_8(z)$
SCDM	0	0.5	1	0	0.5	0.608
SCDM	1	1.4	1	0	0.5	0.304
τ CDM	0	0.5	1	0	0.21	0.608
τ CDM	0.78	1.19	1	0	0.21	0.342
OCDM	0	0.7	0.3	0	0.21	0.85
OCDM	1	1.6	0.46	0	0.21	0.575
Λ CDM	0	0.7	0.3	0.7	0.21	0.9
Λ CDM	1	1.2	0.77	0.23	0.21	0.55

the linear variance $\sigma_L^2(R_0) \equiv 1$ with a Gaussian window function. The function $Q_3(n)$ is given by

$$Q_3(n) = \frac{(4 - 2^n)}{(1 + 2^{n+1})}. \quad (9)$$

Note that at large scales, when the functions $a = b = c = 1$, we recover the tree-level PT expression for the bispectrum; on the other hand, at small scales $a^2 = (7/10) Q_3$ and $b = c = 0$, so Eqs. (2) and (5) give $Q_{123} = Q_3(n)$ independent of triangle configuration: this is the prediction of hyperextended perturbation theory (HEPT), which was shown to describe well the bispectrum (and higher order moments) in the strongly non-linear regime (SF).

Summarizing, the non-linear bispectrum for scale-free initial conditions is given by Eq.(2) and Eqs.(5-8), together with the non-linear power spectrum obtained, for example, from the fitting formula of Peacock & Dodds (1996). In the next section we provide a generalization of the bispectrum fitting formula for CDM models.

3 A FITTING FORMULA FOR CDM MODELS

The numerical simulations used in this work correspond to cluster-normalized cold dark matter (CDM) models run by the Virgo collaboration (see e.g., Jenkins et al. 1998 for more details). There are four different models, SCDM, τ CDM, OCDM, and Λ CDM, see table 1 for a description of the parameters. These simulations each contain 256^3 particles in a box $240h^{-1}$ Mpc on a side. They were run with an adaptive P³M code (Couchman, Thomas & Pearce, 1995; Pearce & Couchman, 1997). There is only one realization for each cosmological model, and all these have the same initial phases. We have also made measurements in a set of smaller simulations run by the Hydra Consortium (Couchman, Thomas & Pearce, 1995), with very similar results. The bispectrum was measured in this set of simulations as described in SCFFHM.

Before generalizing the scale-free fitting formula (SF) reviewed in the previous section, a number of issues related to the CDM spectrum must be addressed. First, since the initial spectrum is no longer scale-free, there is no unique spectral index to be used in Eqs. (6-8). Here we adopt the local spectral index obtained from the linear power spectrum. Another possibility would be to use at scale k the spectral index at scale k_L obtained from the HKLM mapping, $k = k_L [1 + 4\pi k^3 P(k)]^{1/3}$ (Hamilton et al 1991), which maps a non-linear scale to a linear scale with the same number of enclosed pairs. Both approaches are reasonable but can only be




Figure 1. The bispectrum as a function of angle θ between $\mathbf{k}_1$ and $\mathbf{k}_2$ for scales as shown in the panels, for the OCDM model with $\sigma_8 = 0.85$.




Figure 2. The bispectrum as a function of angle θ between $\mathbf{k}_1$ and $\mathbf{k}_2$ for scales as shown in the panels, for the SCDM model with $\sigma_8 = 0.61$.




Figure 3. The bispectrum as a function of angle θ between $\mathbf{k}_1$ and $\mathbf{k}_2$ for scales as shown in the panels, for the Λ CDM model with $\sigma_8 = 0.55$.




Figure 4. The bispectrum as a function of angle θ between $\mathbf{k}_1$ and $\mathbf{k}_2$ for scales as shown in the panels, for the τ CDM model with $\sigma_8 = 0.61$.




Figure 5. The bispectrum for equilateral triangles as a function of scale. Solid lines correspond to the fitting formula, symbols to the measurements in N-body simulations.

simple approximations to a more complex situation; in practice the transfer of power from large to small scales depends not only on the local spectral index but on the full shape of the power spectrum (e.g. Jain & Bertschinger 1994) and non-linear power at a given wavenumber is not just a function of the linear power at another scale. In the transition to the non-linear regime, one-loop calculations (SCFFHM) and numerical simulations with initial power spectra cutoff beyond some characteristic scale (Fry et al. 1993) suggest that the local index approach is a better choice. On the other hand, at small scales one would expect the HKLM choice to be clearly better, since for models such as CDM where the initial power spectrum decreases strongly with k , most of the power is generated by cascading from larger scales. Despite this, approximations for one-point higher order moments in the non-linear regime based on the local spectral index such as EPT (Colombi et al. 1997) and HEPT (SF) show very good agreement even for CDM spectra. Furthermore, attempts to use the HKLM choice gave poor results (Colombi et al. 1997).

Another issue is the dependence of the fitting formula on cosmological parameters such as Ω_m and Ω_Λ . Perturbative results show that the reduced bispectrum Q is very weakly dependent on cosmology at large scales, in particular, $F_2(\mathbf{k}_1, \mathbf{k}_2)$ in Eq. (3) is very well approximated by its Einstein-DeSitter value (Bouchet et al. 1995). In addition, SCFFHM showed that as long as $\Omega_m/f^2(\Omega_m) \approx 1$, where $f(\Omega_m) \approx \Omega_m^{0.6}$, all higher-order moments become independent of cosmology at all orders in perturbation theory for a fixed σ_8 . For these reasons, we shall assume that the fitting formula does not depend on cosmology; this is certainly

a good approximation within the accuracy of our measurements in simulations. Thus, all the cosmology dependence of the bispectrum B_{123} is encoded in the cosmology dependence of the non-linear power spectrum.

We should also note that the asymptotic behavior of the bispectrum at small scales is not yet well tested against N-body simulations. Here we shall follow the standard assumption of the hierarchical ansatz, in which Q becomes constant in the deeply non-linear regime, as implicitly assumed in Eq. (6). This behavior is consistent with our measurements in simulations in the limited range of scales that we can test.

To obtain a fitting formula for the bispectrum in CDM models we proceeded as follows. First, we used the fitting formula for the scale-free case by replacing kR_0 with $q = k/k_{nl}$, where $4\pi k_{nl}^3 P_L(k_{nl}) = 1$, and $P_L(k)$ is the linear power spectrum at the corresponding redshift. This provides a reasonable, but not very accurate, fit to the simulation results, with an average deviation of $d_{\text{avg}} = 40\%$ from the N-body measurements. We then perturbed away from this fit by using two methods. We introduced six free parameters, a_i , such that $q \rightarrow a_1 q$ in Eq. (6), $n + 6 \rightarrow n + a_2$ in Eq. (6), $0.2 \rightarrow 0.2a_3$ in Eq. (7), $4.5 \rightarrow 4.5a_4$ in Eq. (8), $q \rightarrow a_5 q$ in Eq. (8), and finally $[0.7 Q_3(n)]^{1/2} \rightarrow [0.7 Q_3(n)]^{1/2} \sigma_8^{a_6}(z)$, which quantifies deviations from HEPT that depend on redshift. We then obtained the best fit parameters by doing a χ^2 minimization between the predicted and measured bispectrum for all triangles, using a large grid of models (containing about 10^6 of them). We also looked for the best fit using the Levenberg & Marquardt Method (Press et al. 1992), the solution was consistent with the grid approach but some-

what sensitive to the initial guess for a_i . As a result of this procedure we get a best fit model with parameters $a_1 = 0.25$, $a_2 = 3.5$, $a_3 = 2$, $a_4 = 1$, $a_5 = 2$, $a_6 = -0.2$, that is

$$a(n, k) = \frac{1 + \sigma_8^{-0.2}(z) [0.7 Q_3(n)]^{1/2} (q/4)^{n+3.5}}{1 + (q/4)^{n+3.5}} \quad (10)$$

$$b(n, k) = \frac{1 + 0.4 (n+3) q^{n+3}}{1 + q^{n+3.5}} \quad (11)$$

$$c(n, k) = \frac{1 + 4.5 / [1.5 + (n+3)^4] (2q)^{n+3}}{1 + (2q)^{n+3.5}}, \quad (12)$$

and $q = k/k_{nl}(z)$. The average deviation of this fitting formula from the measurements in the simulations is about $d_{\text{avg}} = 15\%$. We also found similarly good fits with $d_{\text{avg}} = 16\%$ for $a_1 = 0.25$, $a_2 = 4$, $a_3 = 1.5$, $a_4 = 1$, $a_5 = 1.5$ and $a_6 = 0$, so the small deviations from HEPT implied by Eq.(10) do not seem statistically significant.

Figures 1-5 illustrate the results. Although we have measured, and used in the fitting procedure, the bispectrum for many triangle shapes, we show for simplicity only triangles where $k_2 = 2k_1$ as a function of angle θ between $\mathbf{k}_1$ and $\mathbf{k}_2$, and equilateral triangles as a function of scale k . Since we only have a single realization for each cosmological model, we cannot show rigorous error bars, which in any case would only be part of the story, since different triangles in a single panel are correlated: highly so in the non-linear regime. However, we can obtain a reasonable estimate of the errors in a single realization by scaling the variance obtained from multiple realizations of Λ CDM lower resolution runs (Scoccimarro et al 2000b): this gives an error in Q of about ± 0.25 for scales $k \gtrsim 0.3$ h/Mpc approximately independent of scale.

Figures 1-4 show a comparison of the fitting formula to the N-body measurements for triangles of different scales (as denoted in each panel) as a function of triangle shape. The reduced bispectrum Q shows the familiar configuration dependence at large scales (top-left panel), and then flattens out at small scales (bottom-right panel). This behavior is well reproduced by the fitting formula, and it is also well understood physically (SCFFHM): at large scales the bispectrum probes the shapes of large-scale structures, which leads to an anisotropic Q , whereas at small scales all configurations become equally probable due to virialization. Note that the flattening is more evident in higher σ_8 models, as expected, since at a given scale these models have a stronger level of non-linearity.

Comparison between corresponding panels in Fig. 3 and Fig. 4 and also between the top panels in Fig. 5 illustrate the lack of sensitivity of Q (for a fixed level of non-linearity, or σ_8) to cosmological parameters such as Ω_m and Ω_Λ , even in the non-linear regime. Note that in some cases the agreement at the smallest scales between the fitting formula and the simulations is not as good as at larger scales, especially in the τ CDM case; however, this does not seem very serious given the estimated error of ± 0.25 and that different triangles are strongly correlated. More numerical work would be necessary to assess this issue.

4 CONCLUSIONS

We provided a fitting formula for the non-linear evolution of the bispectrum in CDM models, which fits the measurements in N-body simulations to an accuracy of 15%. This will be a useful tool for making predictions about the non-linear behavior of non-Gaussian statistics related to the bispectrum of the density field, and is analogous to the role played by fitting formulas for the non-linear evolution of the power spectrum (Hamilton et al. 1991; Jain, Mo & White 1995; Peacock & Dodds 1996). Indeed, Van Waerbeke et al. (2000) apply the results obtained here to the calculation of the skewness of the convergence field in weak gravitational lensing surveys.

There are some issues in this work that merit further investigation. One obvious aspect is improving the accuracy of the fitting function; in view of the upcoming large galaxy and weak lensing surveys it would be useful to have a more accurate prediction for the non-linear evolution of the bispectrum. The main need in this regard is the availability of multiple independent numerical realizations of each cosmological model, as the bispectrum is rather sensitive to the presence of massive clusters in the simulation volume (Colombi et al. 1996). Another issue is the extension towards smaller scales, and the validity of the hierarchical ansatz. Analytical models based on dark matter halos predict that at small scales, $k \gtrsim 3$ h/Mpc, Q should increase with k (Ma & Fry 2000, Scoccimarro et al. 2000b). Testing this prediction reliably will require the use of large-volume, higher-resolution simulations.

5 ACKNOWLEDGMENTS

R.S. is supported by endowment funds from the Institute for Advanced Study and NSF grant PHY-0070928 at IAS. H.M.P.C. Acknowledges support from NSERC. The numerical simulations analyzed in this paper were carried out by the Virgo Supercomputing Consortium (<http://star-www.dur.ac.uk/~frazerp/virgo/virgo.html>) using computers based at the Max Plank Institut fur Astrophysik, Garching and the Edinburgh Parallel Computing Centre.

REFERENCES

- Bernardeau, F. 1994, ApJ, 433, 1
- Bouchet, F. R., Colombi, S., Hivon, E., Juszkiewicz, R. 1995, A & A, 296, 575
- Colombi, S., Bernardeau, F., Bouchet, F. R., & Hernquist, L. 1997, MNRAS, 287, 241
- Colombi, S., Bouchet, F. R., & Hernquist, L. 1996, ApJ, 465, 14
- Couchman, H. M. P., Thomas, P. A., & Pearce, F. R. 1995, ApJ, 452, 797
- Feldman, H., Frieman, J.A., Fry, J.N., Scoccimarro, R. 2000, submitted to Phys. Rev. Lett.
- Frieman, J.A., Gaztañaga, E. 1999, ApJ, 521, L83
- Fry, J. N., & Seldner, M. 1982, ApJ, 259, 474
- Fry, J.N., 1984, ApJ, 279, 499
- Fry, J.N., Melott, A.L., Shandarin, S.F. 1993, ApJ, 412, 504
- Fry, J.N. 1994, Phys. Rev. Lett., 73, 215
- Hamilton, A.J.S., Matthews, A., Kumar, P., Lu, E., 1991, ApJ, 274, L1
- Hivon, E., Bouchet, F.R., Colombi, S., Juszkiewicz, R. 1995, A & A, 298, 643

Jain, B., Bertschinger, E. 1994, ApJ, 431, 495

Jain, B., Mo, H. J., & White, S. D. M. 1995, MNRAS, 276, L25

Jenkins, A. R., Frenk, C. S., Pearce, F. R., Thomas, P. A., Colberg, J. M., White, S. D. M., Couchman, H. M. P., Peacock, J. A., Efstathiou, G. P., & Nelson, A. H. 1998, ApJ 499, 20

Jing, Y.P., Börner, G. 1998, ApJ, 503, 37

Juszkiewicz, R., Bouchet, F. R., & Colombi, S. 1993, ApJ, 412, L9

Ma, C-P., & Fry, J.N. 2000, ApJ, 538, L107

Matarrese, S., Verde, L., & Heavens, A.F. 1997, MNRAS, 290, 651

Peacock, J.A., Dodds, S.J., 1996, MNRAS, 280, L19

Pearce F. R. & Couchman, H. M. P. 1997, New Astronomy, 2, 411

Peebles, P.J.E., 1980, The Large-Scale Structure of the Universe (Princeton: Princeton University Press)

Press, W. H., Teukolsky, S. A., Vetterling, W. T., Flannery, B. P. 1992, Numerical Recipes in Fortran 77, Second Edition, Cambridge University Press, 678

Scoccimarro R., Colombi, S., Fry, J. N., Frieman, J., Hivon, E., & Melott, A., 1998, ApJ, 496, 586 (SCFFHM)

Scoccimarro, R., Frieman, J., 1999, ApJ, 520, 35 (SF)

Scoccimarro, R., Feldman, H., Fry, J.N., Frieman, J.A. 2000, ApJ, in press, astro-ph/0004087

Scoccimarro, R., Sheth, R. K., Hui, L., & Jain, B. 2000b, ApJ, in press, astro-ph/0006319

Van Waerbeke, L., Hamana, T., Scoccimarro, R., Colombi, S., Bernardeau, F. 2000, submitted to MNRAS, astro-ph/0009426